



(12) **Øversettelse av  
europeisk patentskrift**

(11) **NO/EP 2604608 B1**

**NORGE**

(19) NO  
(51) Int Cl.

**C07D 471/04 (2006.01)**

**Patentstyret**


---

- (21) Øversettelse publisert 2016.02.29
- (80) Dato for Den Europeiske Patentmyndighets publisering av det meddelte patentet 2015.10.07
- (86) Europeisk søknadsnr 13151569.4
- (86) Europeisk innleveringsdag 2010.11.22
- (87) Den europeiske søknadens Publiseringsdato 2013.06.19
- (30) Prioritet 2009.11.27, EP, 09177371
- (84) Utpekte stater AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
- (62) Avdelt fra EP2504334, med inndato 2010.11.22
- (73) Innehaver Adverio Pharma GmbH, Willy-Brandt-Platz 2, 12529 Schönefeld, DE-Tyskland
- (72) Oppfinner Mais, Franz-Josef, Otto-Hahn-Strasse 164, 40591 Düsseldorf, DE-Tyskland  
Rehse, Joachim, Am Kloster 12, 42799 Leichlingen, DE-Tyskland  
Joentgen, Winfried, Sprengelstrasse 1, 50735 Köln, DE-Tyskland  
Siegel, Konrad, Augsburgger Strasse 11, 40597 Düsseldorf, DE-Tyskland
- (74) Fullmektig Tandbergs Patentkontor AS, Postboks 1570 Vika, 0118 OSLO, Norge
- 
- (54) Benevnelse **Method for producing methyl- 4,6-diamino-2-[1-(2-fluorobenzyl) -1H-pyrazolo[3,4-b]pyridin-3-yl]-pyrimidin-5-yl-methyl carbamate and its purification for its use as a pharmaceutical agent**
- (56) Anførte publikasjoner WO-A1-03/095451  
WO-A1-2005/046725  
WO-A1-2008/031513  
WO-A1-2010/079120  
CHEMMEDCHEM, Bd. 4, Nr. 5, Mai 2009 (2009-05), Seiten 853-865, XP002622814, ISSN: 1860-7179, DOI: DOI:10.1002/CMDC.200900014  
EVANS R ET AL: "The Preparation of 4-Amino- and Other Pteridines", JOURNAL OF THE CHEMICAL SOCIETY, CHEMICAL SOCIETY, LETCHWORTH; GB, 1. Januar 1956 (1956-01-01), Seiten 4106-4113, XP009144631, ISSN: 0368-1769

- 1 -


**Method for producing methyl-{4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methyl carbamate and its purification for its use as a pharmaceutical agent**

The present invention relates to a process for preparing methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate of the  
5 formula (I)



(I).

Furthermore, a process for purifying the crude product of the compound of the formula  
10 (I) for use as a pharmaceutically active compound, where, for purification, methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate sulphonyldimethane (1:1), i.e. a compound of the formula (II)




- 2 -



(II),

is isolated as intermediate or is generated as intermediate in this purification process, if appropriate present in a mixture, is disclosed

The compound of the formula (I) acts as a stimulator of soluble guanylate cyclase and  
 5 can be used as an agent for the prophylaxis and/or treatment of cardiovascular disorders such as, for example, for the treatment of high blood pressure and of heart failure, stable and unstable angina pectoris, peripheral and cardiac vascular disorders, of arrhythmias, for the treatment of thromboembolic disorders and ischemias such as myocardial infarction, stroke, transitory and ischemic attacks, disturbances of peripheral  
 10 blood flow, prevention of restenoses such as following thrombolysis therapies, percutaneous transluminal angioplasties (PTA), percutaneous transluminal coronary angioplasties (PTCA), bypass and for the treatment of arteriosclerosis, asthmatic disorders and diseases of the urogenital system such as, for example, prostate hypertrophy, erectile dysfunction, female sexual dysfunction, osteoporosis, glaucoma, pulmonary  
 15 hypertension, gastroparesis and incontinence.

The preparation of the compound of the formula (I) and its purification are known in principle. WO 03/095451 describes the preparation of the compound of the formula (I) by the route below.





5

Here, initially 2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-[(E)-phenyldiazenyl]pyrimidine-4,6-diamine of the formula (III) is cleaved by catalytic hydrogenation, and the resulting trisamino compound is isolated as 2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-4,5,6-pyrimidinetriamine

10 trihydrochloride of the formula (IV). This trihydrochloride is then reacted with methyl chloroformate of the formula (V) in the solvent pyridine to give methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinylcarbamate of the formula (VI). Alternatively, ChemMedChem 2009, 4, 853-865 describes that the trisamino compound is isolated as trihydrochloride and the HCl-free base is then generated by extraction with aqueous NaHCO<sub>3</sub> solution and the free base is reacted

15 with methyl chloroformate of the formula (V) in the solvent pyridine to give the

- 4 -

compound of the formula (VI). The compound of the formula (VI) is then reacted with methyl iodide of the formula (VII) in the presence of a base to give the crude product of the compound of the formula (I). The crude product of the compound of the formula (I) is purified according to the experimental procedure of Example 8 of WO 03/095451 and the comparable description in ChemMedChem 2009, 4, 853-865 by triturating the crude product with dichloromethane/THF, intermediate isolation of the product triturated with dichloromethane/THF by filtration, boiling the isolated solid with methanol, intermediate isolation of the solid boiled with methanol by filtration, dissolution of the solid in a mixture of dioxane, dichloromethane and methanol in the presence of activated carbon, removal of the activated carbon by filtration through kieselguhr or Celite, concentration of the filtered solution to dryness, trituration of the solid concentrated to dryness with methanol, isolation of the solid triturated with methanol by filtration and (not described in WO 03/095451 in Example 8 or ChemMedChem 2009, 4, 853-865, but objectively required) drying. Alternatively, a crude product of the compound of the formula (I) concentrated to dryness can be purified in poor yields by preparative chromatography (RP-HPLC).

This synthesis and the purifications have a number of disadvantages which are very unfavourable for an industrial realization on a large scale. This is true especially for the isolation of the trisamino compound as trihydrochloride of the formula (IV). The addition of hydrochloric acid requires an acid-proof industrial plant, and the yield of the step is only an unsatisfactory 59.3% of theory (see, for example, Example 8A of WO 03/095451). The realization of the reaction of the trisamino compound of the formula (IV) or the corresponding HCl-free base in the solvent pyridine is likewise disadvantageous. The compound of the formula (VI) can only be isolated by complete evaporation of the reaction mixture, which is disadvantageous on an industrial scale (see, for example, Example 5 of WO 03/095451). On a relatively large scale, such steps generally result in considerable problems such as sticking-on or thermal decomposition owing to the substantially longer thermal stress when a reaction is carried out on a relatively large scale. The purification of the product of the formula (VI) according to the experimental procedure of Example 5 from WO 03/095451 by boiling in diethyl ether, too, has considerable disadvantages. Because of the high


- 5 -

flammability of diethyl ether, this step can be realized only with increased industrial expenditure.

However, particularly disadvantageous are the purification processes for the crude product of the formula (I). An effective purification is a *conditio sine qua non* for use  
5 as a pharmaceutically active compound. The described purification via RP HPLC, i.e. the chromatographic purification, is a laboratory method, the realization of which on an industrial scale is very expensive. In addition, the stated yield of only 29% for the synthesis step to the crude product of the formula (I) and its purification is very low.  
10 The alternative preparation and purification method is very complicated. It comprises a total of 5 isolations of solids (2 concentrations to dryness and 3 filtrations), and, as already mentioned above, concentrations to dryness on an industrial scale are very unfavourable. Altogether, when carrying out a chemical step, a number of 5 isolations of solids for the preparation and purification of a pharmaceutically active compound on an industrial scale is very disadvantageous.

15 Accordingly, it was the object to provide a simplified process which is safe and can also be carried out advantageously on an industrial scale and which affords an active compound in high yield and high purity in pharmaceutically acceptable quality.

Surprisingly, we have now found a process for preparing methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate of the  
20 formula (I)




(I)

- 6 -


and its purification for use as a pharmaceutically active compound. This novel process and the purification of the crude product of the compound of the formula (I) differ from the processes known to date in the following points:

- 5
- After catalytic hydrogenation of the compound of the formula (III), the trisamino compound is isolated as the free base of the formula (VIII) without intermediate formation of salts



(VIII).


- 10
- The preparation of the compound of the formula (VI) is carried out using methyl chloroformate or dimethyl dicarbonate as reagent in a pyridine-free process



(VI).

- 7 -

- 5 - The compound of the formula (VI) is converted in a manner known per se using a methylating agent into a crude product of the formula (I); the purification of the crude product of the formula (I) for use as pharmaceutically active compound is carried out via the compound methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate sulphinyldimethane (1:1), i.e. a compound of the formula (II) as isolated intermediate or generated in a mixture



(II).


- 10 By virtue of these differences, it is possible to overcome the disadvantages of the processes known to date and to obtain an active compound in high yield and high purity and pharmaceutically acceptable quality.

The process for preparing the compound of the formula (I) and the purification via the intermediate of the formula (II) are described in detail below.

### 15 **Catalytic hydrogenation of the compound of the formula (III)**

The first step of the process according to the invention begins with a catalytic hydrogenation of the compound of the formula (III).

- 8 -



(free base of the formula  
(IV))

- 5 This may be carried out in the presence of Raney nickel or industrially customary Pt/carbon or Pd/carbon catalysts. Preference is given to Pt/carbon and Pd/carbon. N,N-Dimethylformamide (DMF), N,N-dimethylacetamide (DMA) or N-methyl-2-pyrrolidone (NMP), preferably DMF, serves as solvent.

10 Hydrogenation conditions are temperature 40-80°C, preferably 50-70°C, pressure: 2-90 bar, preferably 5-70 bar, of hydrogen, hydrogenation time: 1-72 h, preferably 3-36 h.

After removal of the catalyst by filtration, the product is precipitated from a C<sub>1</sub>-C<sub>4</sub>-alcohol, preferably methanol or ethanol, and/or water. Preference is given to a mixture of methanol, isopropanol or ethanol and water.

- 15 In the context of the invention, a C<sub>1</sub>-C<sub>4</sub>-alcohol is a straight-chain or branched alcohol having 1 to 4 carbon atoms. The following may be mentioned by way of example and by way of preference: methanol, ethanol, n-propanol, isopropanol, n-butanol and tert-butanol. This definition also applies to the C<sub>1</sub>-C<sub>4</sub>-alcohols used hereinbelow.


20 It is also possible to remove some of the solvent prior to the precipitation, a partial distillative removal of 0-80%, preferably 40-70%, of the solvent is in accordance with the invention.

- 9 -

The moist product obtained in this manner is dried under reduced pressure: this gives the product of the formula (VIII) (corresponds to the free base of the formula (IV)).

**Reaction of the compound of the formula (VIII) with methyl chloroformate (V)**

It is furthermore disclosed that the product of the formula (VIII) is then reacted, for example, with methyl chloroformate of the formula (V) in a novel pyridine-free process to give the product of the formula (VI).




The solvent used for the reaction is a C<sub>1</sub>-C<sub>4</sub>-alcohol, preferably ethanol, methanol, isopropanol, particularly preferably isopropanol.

The amount of methyl chloroformate is from 1.0 to 3.0 equivalents, preferably from 1.0 to 2.0 equivalents, based on the compound of the formula (VIII) employed.

Possible reaction temperatures are 0-75°C, preferably 15-50°C.

During the reaction, hydrogen chloride is formed which forms a compound of the formula (IX), i.e. the hydrochloride of the product of the formula (VI), in the reaction mixture. This product of the formula (IX) can either be isolated as HCl-containing product and be cleaved by addition of base to the product of the formula (VI), or it can be cleaved by addition of base even before the isolation, so that the product of the formula (VI) is isolated directly.



It is preferred to cleave the product of the formula (IX) by addition of the base prior to the isolation and to isolate the free base of the formula (VI) directly.

- 5 Suitable bases are all bases having a pKB which is higher than that of the compound of the formula (I). Examples which may be mentioned are: hydroxides, carbonates and phosphates of the alkali metals and alkaline earth metals, nitrogen-containing organic bases, such as trialkylamines, guanidines or amidines. Examples which may be mentioned are: lithium hydroxide, sodium hydroxide, potassium hydroxide, rubidium
- 10 hydroxide, caesium hydroxide, magnesium hydroxide, calcium hydroxide, strontium hydroxide, barium hydroxide, lithium carbonate, sodium carbonate, potassium carbonate, rubidium carbonate, caesium carbonate, magnesium carbonate, calcium carbonate, strontium carbonate and barium carbonate, sodium phosphate and potassium phosphate, trialkylamines having straight-chain, cyclic or branched C<sub>1</sub>-C<sub>20</sub>-
- 15 alkyl radicals, and cyclic or open-chain guanidines or amidines. Preference is given to triethylamine, tripropylamine, diisopropylethylamine, tributylamine, dicyclohexylethylamine, cyclohexyldimethylamine, cyclohexyldiethylamine, triisooctylamine, tridecylamine, tridodecylamine, trihexadecylamine, N-methylmorpholine, DBU, DBN, tetramethylguanidine, etc. Particular preference is
- 20 given to triethylamine, tributylamine, diisopropylethylamine, N-methylmorpholine, DBU, DBN.

The amount of base is from 1.0 to 2.0 equivalents, preferably from 1.0 to 1.5 equivalents, based on the methyl chloroformate of the formula (V) employed.


- 11 -

Possible reaction temperatures for the reaction with the base are 0-100°C, preferably 15-70°C.

The product of the formula (VI) is present in suspension and is isolated by filtration. It is washed with the C<sub>1</sub>-C<sub>4</sub>-alcohol and dried under reduced pressure in a customary manner.

### Reaction of the compound of the formula (VIII) with dimethyl dicarbonate (X)

In a further process disclosed, the product of the formula (VIII) is reacted with dimethyl dicarbonate of the formula (X) to give the product of the formula (VI). This reaction does not require any base such as, for example, pyridine.



The solvents used for this reaction are C<sub>1</sub>-C<sub>4</sub>-alcohols, preferably ethanol, methanol, isopropanol, particularly preferably isopropanol.

The amount of dimethyl dicarbonate is from 1.0 to 3.0 equivalents, preferably from 1.0 to 2.0 equivalents, based on the compound of the formula (VIII) employed.

Possible reaction temperatures are 0-65°C, preferably 15-40°C.

The product of the formula (VI) precipitates and is isolated by filtration. It is washed with the C<sub>1</sub>-C<sub>4</sub>-alcohol and dried under reduced pressure in a customary manner.

- 12 -

In the reaction with dimethyl dicarbonate, the product of the formula (VI) is obtained directly. Further addition of base is therefore not required.

Both processes, i.e. the reaction of the compound of the formula (VIII) with methyl chloroformate and subsequent cleavage of the hydrochloride of the formula (IX) with  
5 base or the reaction of the compound of the formula (VIII) with dimethyl dicarbonate afford a comparable quality of the product of the formula (VI), so that the products of the formula (VI) from the two processes can be used in the same manner for the further conversion into the product of the formula (I).


Both processes are preferred.

10 The compound of the formula (VI) can form solvates or solvent-containing solid forms, for example methanol-, ethanol-, or isopropanol-containing solid forms. It is therefore possible, that, when the hydrochloride of the formula (IX) is cleaved to the product of the formula (VI) or when the product of the formula (VI) is synthesized directly with dimethyl dicarbonate, a solvate of the C<sub>1</sub>-C<sub>4</sub>-alcohol used as solvent is obtained. The  
15 solvate may be so stable that, during drying of the product of the formula (VI), it does not decompose completely, and clearly noticeable solvent residues, i.e., for example, of the C<sub>1</sub>-C<sub>4</sub>-alcohol in question, thus remain in the product of the formula (VI). On the other hand, the product of the formula (VI) must not be dried at temperatures which are too hot, since it may decompose with formation of byproducts at  
20 temperatures which are too high.

Accordingly, it is preferred to dry the product of the formula (VI) from the cleavage of the hydrochloride of the formula (IX) with base or from the direct synthesis with dimethyl dicarbonate at a product temperature of not more than 110°, particularly preferably at a product temperature of not more than 100°. Here, it is particularly  
25 preferred for any residues of C<sub>1</sub>-C<sub>4</sub>-alcohol present as solvate to remain in the product of the formula (VI) and to use the product of the formula (VI) in this form for preparing the intermediate of the formula (II) or the product of the formula (I). With very particular preference, the product of the formula (VI) contains isopropanol as residual solvent in a range of from 0 to 13%.

### Methylation of the compound of the formula (VI)


The product of the formula (VI) obtained in this manner is reacted in a manner known per se, for example in accordance with one of the descriptions in WO 03/0945451 or ChemMedChem 2009, 4, 853-865, with a methylating agent Me-X to give a crude product which contains high amounts of the compound of the formula (I).



The methylating agent Me-X used is methyl iodide, dimethyl sulphate, methyl toluenesulphonate, etc., and methyl iodide or dimethyl sulphate is preferred.

### 10 Purification of the crude product of the compound of the formula (I)

The crude product of the formula (I) is purified for use as pharmaceutically active compound. To this end, initially, a mixture is formed which contains high amounts of the compound of the formula (II) as intermediate.



- 14 -

(I) (crude product)

(II)

To this end, the crude product of the formula (I) is dissolved in DMSO, if appropriate in the presence of a pharmaceutically acceptable simple solvent from the class of the ketones, ethers, esters or alcohols. Examples of such solvents which may be mentioned  
5 are: methanol, ethanol, isopropanol, 1-butanol, 2-butanol, ethyl acetate, isopropyl acetate or propyl acetate, butyl acetate, tert-butyl methyl ether, diisopropyl ether, acetone, methyl ethyl ketone, methyl isobutyl ketone, etc. Preference is given to ethanol, isopropanol, ethyl acetate, isopropyl acetate, butyl acetate, methyl ethyl ketone, methyl isobutyl ketone; particular preference is given to ethyl acetate. It is also  
10 possible to use mixtures of these solvents.

DMSO is added in an amount of from 100 to 750% by weight, preferably from 150 to 500% by weight, based on the amount of the crude product of the formula (I) employed.

If appropriate, activated carbon may be added to this mixture in an amount of from  
15 0.25 to 35% by weight, preferably from 0.5 to 20% by weight, based on the amount of the crude product of the formula (I) employed.

To form a solution, the mixture is heated to 40-120°C, preferably 50-100°C.

To form a pharmaceutically acceptable product of the formula (I), the solution has to be filtered. The filtration has to be carried out independently of whether activated  
20 carbon was added or not.

The amount of the pharmaceutically acceptable solvent which, in addition to DMSO, is added to the solution of the crude product of the formula (I), i.e. used prior to the filtration, is from 25 to 200% by weight, preferably from 40 to 100% by weight, based on the DMSO.

25 The filtration is carried out hot, the temperatures are 40-120°C, preferably 50-100°C.

After the filtration, a pharmaceutically acceptable solvent, preferably the same solvent as above, is added to the hot filtrate. This results in a crystallization of the product of the formula (II).

- 15 -

The total amount of solvent added before and after the filtration is from 200 to 1500% by weight, preferably 400-1200% by weight, based on the DMSO.

The addition temperature is 30-110°C, preferably 35-90°C.

5 Prior to the isolation of the solid which contains high amounts of the compound of the formula (II), to bring the precipitation to completion, the mixture is cooled to a temperature range of 0-35°C, preferably to an ambient temperature of, for example, 20-30°C.

10 The isolation is carried out using customary isolation devices such as a nutsche filter or a centrifuge. To remove the mother liquor, the isolated material is, during isolation, washed with a pharmaceutically acceptable solvent, the same solvent as above being preferred.


15 After the DMSO redissolution, the isolated material contains high amounts of the product of the formula (II). In addition, small amounts of the product of the formula (I) may also usually precipitate directly without forming a solvate with DMSO. Also possible is the formation of solvates of a different stoichiometry or the formation of solvent adducts with no fixed stoichiometry. Moreover, DMSO may also be present in unbound form as an adhering residual solvent. The content of DMSO in the isolated material is usually from 10 to 25% by weight, preferably 12-17%. The product of the formula (II) is particularly preferably formed in the form of this mixture and used for  
20 preparing the purified product of the formula (I).

The product of the formula (II) obtained in this manner can now be dried or, alternatively, be used in moist form comprising solvent residues, i.e. adhering DMSO and the precipitation solvent(s), for conversion into the purified product of the formula (I).

25 The compound of the formula (II) is novel. It can be prepared in pure form as described in the working examples below and be characterized analytically.

For pharmaceutical use, the DMSO has to be removed from the product of the formula (II) or the mixture comprising high amounts of the compound of the formula (II).

- 16 -



To this end, the product of the formula (II) or the isolated mixture comprising high amounts of the product of the formula (II) is boiled in a pharmaceutically acceptable solvent from the class of the ketones, ethers, esters or alcohols. Examples of such solvents which may be mentioned are: methanol, ethanol, isopropanol, 1-butanol, 2-butanol, ethyl acetate, isopropyl acetate or propyl acetate, butyl acetate, tert-butyl methyl ether, diisopropyl ether, acetone, methyl ethyl ketones, methyl isobutyl ketone, etc. Preference is given to ethanol, isopropanol, ethyl acetate, isopropyl acetate, butyl acetate, methyl ethyl ketone, methyl isobutyl ketone. It is also possible to use mixtures of these solvents. Particular preference is given to ethyl acetate or a mixture of ethyl acetate with ethanol.

Boiling takes place at reflux of the solvent in question or, if appropriate, at slightly elevated pressure. The temperature is 50-150°C, preferably 80-120°C.

The process disclosed offers marked advantages compared to the prior art. Surprising was in particular that the direct isolation of the compound of the formula (VIII) (free base) without intermediate formation of the compound of the formula (IV) (trihydrochloride) allowed the yield to be increased markedly, with a simultaneous markedly more simple industrial practice (no acid-proof parts of the plant).

The compound of the formula (VIII) can then be converted in novel pyridine-free processes with methyl chloroformate or dimethyl carbonate into the compound of the formula (VI). These novel processes are very simple and can be carried out with minimum expense in industry. After the reaction, the product of the formula (VI) is

- 17 -

present suspended as a solid and can be isolated without evaporation steps by filtration. The yield obtained is very high.

It is furthermore disclosed that the purification of the crude product of the formula (I) for pharmaceutical use takes place in particular by redissolution in a DMSO-containing solvent mixture and that the novel compound of the formula (II) is obtained as an intermediate in this step, if appropriate in a mixture in high proportions. By this step, all impurities are removed except for small residual amounts, so that, after the DMSO content has been removed by simple boiling, a highly pure solid of the formula (I) remains. This solid is generally colourless to very slightly yellow and the analytical purity (HPLC) is markedly above 98% by weight, which is very advantageous for pharmaceutical use.

The process can be carried out safely technically and allows a production on an industrial scale. It can be adapted flexibly to existing apparatus in the plant. In a particularly preferred embodiment, in the purification of the crude product of the formula (I), the intermediate isolation of the product of the formula (II) or of the mixture comprising high amounts of the compound of the formula (II) is carried out in a nutsche filter dryer. Subsequent removal of the DMSO from the product of the formula (II) isolated as an intermediate in the nutsche filter dryer is carried out by direct addition of solvent to the nutsche filter dryer with or without intermediate drying of the product of the formula (II). This avoids open handling of the solid of the product of the formula (II) with the associated risk of contamination.

**Experimental part****Abbreviations and acronyms:**

abs.	absolute
cat.	catalytic
CI	chemical ionization (in MS)
D	day(s)
TLC	thin-layer chromatography
DMF	dimethylformamide
DMSO	dimethyl sulphoxide
Ee	enantiomeric excess
EI	electron-impact ionization (in MS)
Ent	enantiomer/enantiomerically pure
Eq	equivalent(s)
ESI	electrospray ionization (in MS)
GC-MS	gas chromatography-coupled mass spectrometry
% by weight	per cent by weight
H	hour(s)
HPLC	high-pressure, high-performance liquid chromatography
conc.	concentrate
LC-MS	liquid chromatography-coupled mass spectrometry

- 19 -

Min	minute(s)
MS	mass spectrometry
NMR	nuclear magnetic resonance spectrometry
Ph	phenyl
R <sub>f</sub>	retention index (in TLC)
R <sub>t</sub>	retention time (in HPLC)
RT	room temperature
v/v	volume-to-volume ratio (of a solution)
aq.	aqueous, aqueous solution

**Examples 1 and 3 below illustrate the invention, but the invention is not limited to the examples.**

### **Example 1**

#### **Preparation of 2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-4,5,6-pyrimidinetriamine (VIII)**

In a pressure autoclave, 1100 g of the compound of the formula (III) were suspended in 5.4 l of DMF. 44 g of a conventional water-moist (about 50%) 5% Pd/carbon catalyst were added, and the sealed autoclave was, after inertization with nitrogen and application of hydrogen, hydrogenated at a hydrogen pressure of 65 bar and an internal  
10 temperature of about 60°C for about 18 h. After cooling to about 25°C, venting and inertization, the autoclave content was removed, rinsing with 650 ml of DMF.

Three of such reactions carried out in the same manner were combined, the old catalyst was filtered off, the filtercake was rinsed with 1.1 l of DMF and the filtrate was concentrated under reduced pressure to about one third of its mass. Successively, 8.25 l  
15 of methanol and 8.25 l of water were metered into the residue of about 6.5 kg, to bring the crystallization to completion, the suspension was cooled to about 5°C and the solid was filtered off and washed with methanol/water (1:1 vol). The product was dried at 50°C under reduced pressure. The weight was 2415 g, which corresponds to 91.8% of theory. The content of the target product of the formula (VIII) (free base) was >98  
20 area% or >97% by weight. The most significant impurities were DMF (about 0.8% by weight) and water (about 0.5% by weight).

### **Disclosed and unclaimed Example 2**

#### **Preparation of methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinylcarbamate (VI)**

25 3063 g of the compound of the formula (VIII) and 30.7 l of technical grade isopropanol were initially charged in a reaction vessel. With stirring, 1641 g of dimethyl dicarbonate were metered in at 20-25°C, and the mixture was stirred at this temperature for 22 h. The precipitated product was filtered off with suction, washed

- 21 -

with industrial grade isopropanol and dried at 50°C under reduced pressure. The weight of the product obtained was 3748 g or 105.9% of theory. The product of the formula (I) contained, inter alia, about 4.7% of isopropanol virtually unremovable by drying (partially, an isopropanol solvate was present), and the analytical content was 89.5% by weight (HPLC). Based on this content, the yield was 94.8% of theory.

### **Disclosed and unclaimed Example 3**

#### **Preparation of 2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-4,5,6-pyrimidinetriamine (VIII)**

In a pressure autoclave, 300 g of the compound of the formula (III), 1600 ml of DMF and 60 g of water-moist Raney nickel were initially charged and, after inertization, hydrogenated at an internal temperature of 60°C and a hydrogen pressure of 65 bar for about 18 h. After cooling and venting, the old catalyst was filtered off and rinsed with 100 ml of DMF. The filtrate was concentrated under reduced pressure to 534.5 g, and at 35-40°C, 750 ml of methanol and then, after cooling, at 0-5°C, 750 ml of water were metered into the residue. The solid was filtered off and dried at 50°C under reduced pressure. The weight was 219.7 g or 91.8% of theory.

### **Disclosed and unclaimed Example 4**

#### **Preparation of methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinylcarbamate (VI)**

In a reaction vessel, 1.50 kg of the compound of the formula (VIII) were initially charged in 14.25 l of isopropanol, and the mixture was heated with stirring to 35°C. 531 g of methyl chloroformate were, at a steady rate, metered in over a period of 30 min, rinsing with 750 ml of isopropanol, and the mixture was stirred at 35°C for 16 h. The mixture was then heated to 50°C and 3.85 l of methanol and 606 g of triethylamine were metered in with stirring at 50°C, rinsing with 450 ml of methanol. The mixture was then stirred at 50°C for 1 h, cooled to RT and stirred at RT for 1 h. The suspended solid was filtered off with suction, washed twice with in each case 3.0 l of isopropanol/methanol (4:1) and once with 3.0 l of isopropanol and sucked dry. The

- 22 -

moist product was dried at 50°C for 1 h and then at 100°C for 22 h in a vacuum drying cabinet. The weight of the product obtained was 1.793 kg or 103.3% of theory. The product of the formula (VI) contained 6.45% of isopropanol virtually unremovable by drying (partially, an isopropanol solvate was present), and the analytical content was 87.9% by weight (HPLC). Based on this content, the yield was 90.8% of theory.

#### **Disclosed and unclaimed comparative example 5**

##### **Preparation of methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinyl(methyl)carbamate (I)**

(Methylation in a manner known per se in accordance with WO 03/095451, Example 8 second procedure)

At 20-25°C, 1630 g of the compound of the formula (VI) was suspended in 16.3 l of THF. The suspension was cooled to from -6 to -4°C, and 3480 g of a 1M solution of bis(trimethylsilyl)sodium amide were metered in. The mixture was stirred, 596 g of methyl iodide were metered in, the mixture was stirred briefly and slowly allowed to warm to about 5°C. The mixture was stirred at this temperature until the reaction had ended (about 4 h). The reaction mixture was washed 4 times with 4.1 l of 15% strength ammonium chloride solution. The organic phase was concentrated by evaporation to a residue of about 6.4 kg, and the temperature was adjusted to about 25°C. The precipitated solid was filtered off, washed with a total of 3 l of THF and dried at 50°C under reduced pressure. This gave 1112 g of the crude product of the formula (I). This corresponded to a yield of 75.2% of theory.

#### **Disclosed and unclaimed Example 6**

**Preparation of a mixture consisting of methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinyl(methyl)carbamate (I) and methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate sulphinyldimethane (II) with high amounts of the product of the formula (II)**

- 23 -

9.0 g of a crude product of the formula (I) which had been prepared in a manner comparable to Comparative Example 5 were dissolved in 16 ml of DMSO at 100°C. (The clarification by filtration which would have been required at this point to achieve a pharmaceutically acceptable product quality was dispensed within this laboratory experiment). The mixture was then allowed to cool to 75°C, 110 ml of ethyl acetate were added and the mixture was cooled slowly to about 25°C. The precipitated solid was filtered off, washed with a total of 28 ml of ethyl acetate and dried at 50°C under reduced pressure. The weight was 9.6 g or 90.0% of theory.

#### **Disclosed and unclaimed Example 7**

#### **10 Preparation of purified methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinyl(methyl)carbamate (I)**

The entire amount of the product of the formula (II) prepared in the above Example 6 was stirred in 135 ml of ethyl acetate at reflux (about 78°C) for 1 h and cooled to about 25°C. The solid was filtered off with suction, washed with a total of 36 ml of ethyl acetate and dried under reduced pressure. The weight was 7.6 g or 93.8% of theory. The content of the product was markedly above 98% by weight (HPLC). As solvent, ethyl acetate was present in an amount of about 0.2%. The DMSO content was below 0.1%.

#### **Disclosed and unclaimed Example 8**

#### **20 Preparation of purified methyl 4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-pyrimidinyl(methyl)carbamate (I) with intermediate isolation of a mixture comprising high amounts of methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate sulphinyldimethane (II) as moist product**

25 193.5 g of a crude product of the formula (I) which had been prepared in a manner comparable to Comparative Example 5 were dissolved in 344 ml of DMSO and 172 ml of ethyl acetate at about 96°C. 19.4 g of activated carbon and 172 ml of ethyl acetate were then added, and the hot mixture was stirred. The hot mixture was then filtered off

- 24 -

to remove the activated carbon, rinsing with 172 ml of ethyl acetate. The temperature of the filtrate was adjusted to 78°C, and 1850 ml of ethyl acetate were added slowly. Over about 2-3 h, the mixture was cooled to about 25°C, and the solid was filtered off and washed with a total of 772 ml of ethyl acetate. The moist product, which contained high amounts of the compound of the formula (II) in a mixture was suspended in 2900 ml of ethyl acetate, heated at reflux for 1 h and cooled to about 25°C. The solid was filtered off with suction, washed with a total of 774 ml of ethyl acetate and dried at 50°C under reduced pressure. The weight obtained was 155.1 g or 80.2% of the starting material. The content of the product was markedly above 98% by weight (HPLC). As solvents, virtually only ethyl acetate and DMSO were present in small amounts.

#### **Disclosed and unclaimed Example 9**

**Preparation and analytical characterization of methyl {4,6-diamino-2-[1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}methylcarbamate sulphinyldimethane (II)**

14.8 g of a crude product of the formula (I) which had been prepared in a manner comparable to Comparative Example 5 were dissolved in 28.9 g of DMSO and 11.85 g of ethyl acetate at about 94°C. 1.5 g of activated carbon Norit A-Supra and a further 11.85 g of ethyl acetate were then added, the mixture was stirred at reflux (88-90°C) for 1 h and the hot mixture was then filtered to remove the activated carbon. The solid, some of which had already precipitated, was re-dissolved by warming to about 78°C, and the solution was then allowed to cool slowly. The precipitated solid was filtered off with suction at RT, washed three times with in each case 50 ml of ethyl acetate and dried in a drying cabinet at 30°C for 18 h. This gave 9.2 g or 52.5% of theory of a slightly yellowish crystal powder of the compound of the formula (II).

HPLC: 99.90 area% (without taking the DMSO into account)

DMSO (GC): 14.7% by weight

<sup>1</sup>H-NMR (400 MHz in DMF-d<sub>7</sub>):

- 25 -

d = 2.59 (s, about 6H, 2 CH<sub>3</sub> at DMSO), 3.13 (s, 3H, N-CH<sub>3</sub>), 3.58 + 3.67 (two s, 3H, hindered rotation at O-CH<sub>3</sub>), 5.91 (s, 2H, -CH<sub>2</sub>-), 6.53 (s, 4H, 2 -NH<sub>2</sub>), 7.05-7.40 (m, 5H, 4 aromatic H at the o-fluorobenzyl substituent and 1H at the pyrido ring meta to the pyrido nitrogen), 8.60 (dd, 1H, at the pyrido ring ortho to the pyrido nitrogen), 9.12  
5 (dd, 1H, at the pyrido ring para to the pyrido nitrogen).

Elemental analysis:

found	C: 52.2%	calculated	C: 52.79%
	H: 4.9%		H: 5.03%
	N: 22.7%		N: 22.39%

## Patentkrav

- 5 **1.** Fremgangsmåte for fremstilling av metyl{4,6-diamino-2-[1-(2-fluorbenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}metylkarbamat med formel (I) **karakterisert ved at** 2-[1-(2-fluorbenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-[(E)-fenyldiazenyl]pyrimidin-4,6-diamin med formel (III) spaltes ved katalytisk hydrogenering og **i det** 2-[1-(2-fluorbenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-4,5,6-pyrimidintriamin med formel (VIII), etter filtrering av katalysatoren, utfelt fra en
- 10 C<sub>1</sub>-C<sub>4</sub>-alkohol og/eller vann, uten mellomliggende dannelse av salter.
- 2.** Fremgangsmåte ifølge krav 1 for fremstilling av metyl{4,6-diamino-2-[1-(2-fluorbenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]pyrimidin-5-yl}metylkarbamat med formel (I), **karakterisert ved at** 2-[1-(2-fluorbenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-5-[(E)-fenyldiazenyl]pyrimidin-4,6-diamin med formel (III) spaltes ved katalytisk
- 15 hydrogenering og **i det** 2-[1-(2-fluorbenzyl)-1H-pyrazolo[3,4-b]pyridin-3-yl]-4,5,6-pyrimidintriamin med formel (VIII), etter filtrering av katalysatoren, utfelt fra metanol og/eller vann, uten mellomliggende dannelse av salter.