

### (12) Oversettelse av europeisk patentskrift

NORGE	(19) NO (51) Int Cl.
	A61K 9/19 (2006.01)
	A61K 9/00 (2006.01)
	A61K 9/08 (2006.01)
	A61K 9/16 (2006.01)
	A61K 31/00 (2006.01)
	A61K 31/407 (2006.01)
	A61K 47/10 (2017.01)
	A61K 47/18 (2017.01)
	A61K 47/26 (2006.01)

### Patentstyret

(45)	Oversettelse publ	lisert	2020.02.17
(80)	Dato for Den Euro Patentmyndighets publisering av det patentet	opeiske s t meddelte	2019.11.20
(86)	Europeisk søknad	dsnr	17188743.3
(86)	Europeisk innleve	eringsdag	2013.10.22
(87)	Den europeiske s Publiseringsdato	øknadens	2018.01.24
(84)	Utpekte stater		AL ; AT ; BE ; BG ; CH ; CY ; CZ ; DE ; DK ; EE ; ES ; FI ; FR ; GB ; GR ; HR ; HU ; IE ; IS ; IT ; LI ; LT ; LU ; LV ; MC ; MK ; MT ; NL ; NO ; PL ; PT ; RO ; RS ; SE ; SI ; SK ; SM ; TR
(62)	Avdelt fra		EP2865391, 2013.10.22
(73)	Innehaver		medac Gesellschaft für klinische Spezialpräparate mbH, Theaterstrasse 6, 22880 Wedel, Tyskland
(72)	Oppfinner		Schuldt-Lieb, Sonja, Charlottenstraße 26, 20257 Hamburg, Tyskland Bialleck, Sebastian, Guldtweg 1, 22587 Hamburg, Tyskland Guhde, Ingo, Godeffroystraße 30, 22587 Hamburg, Tyskland Rehberg, Michaela, Sülldorfer Kirchenweg 72 a, 22587 Hamburg, Tyskland
(74)	Fullmektig		ZACCO NORWAY AS, Postboks 2003 Vika, 0125 OSLO, Norge
(54)	Benevnelse	METHOD F	OR PRODUCING A FREEZE-DRIED PHARMACEUTICAL COMPOSITION
(56)	Anførte publikasjoner	US-A1- 200 GB-A- 2 17	09 162 440 1 408

Vedlagt foreligger en oversettelse av patentkravene til norsk. I hht patentloven § 66i gjelder patentvernet i Norge bare så langt som det er samsvar mellom oversettelsen og teksten på behandlingsspråket. I saker om gyldighet av patentet skal kun teksten på behandlingsspråket legges til grunn for avgjørelsen. Patentdokument utgitt av EPO er tilgjengelig via Espacenet (<u>http://worldwide.espacenet.com</u>), eller via søkemotoren på vår hjemmeside her: <u>https://search.patentstyret.no/</u>

#### Description

5

The invention relates to a freeze-dried pharmaceutical composition comprising mitomycin C, which is characterized by high stability and purity and can rapidly be reconstituted to form ready-to-use solutions.

Mitomycin C, chemical name (1aS,8S,8aR,8bS)-6-amino-8-[[(aminocarbonyl)oxy]methyl]-1,1a,2,8,8a,8b-hexahydro-8a-meth-oxy-5-methyl-azirino[2',3':3,4]pyrrolo[1,2-a]indol-4,7dione or 6-amino-8a-methoxy-5-methyl-4,7-dioxo-1,1a,2,4,7,8,8a,8b-

- 10 octahydroazarino[2',3':3,4]pyrrolo[1,2-a]indol-8-ylmethyl carba-mate, was first isolated in 1958 from Streptomyces caespitosus (see Wakaki et al., Antibiotics and Chemotherapy, 8, 228, 1958 and GB 830874). It has excellent antitumour activity and can above all be used for the treatment of bladder tumours, gastric carcinomas, bronchial carcinomas, pancreatic carcinomas, colorectal carcinomas, mammary carcinomas, liver cell carcinomas, cervical carcinomas, oesophageal
- 15 carcinomas, carcinomas in the head and neck region, chronic myeloid leukaemia and osteosarcoma. For this, it is used in the form of aqueous solutions for injection or infusion or solutions for intravesicular use.

US 5,216,011 describes solutions of mitomycin C which can for example be administered by injection. These solutions are intended to circumvent difficulties in the preparation of freezedried preparations. As solvent, the solutions comprise 40 to 100 vol.-% propylene glycol and 0 to 60 vol.-% water.

US 2009/0162440 describes freeze-dried microspheres containing mitomycin C.

25

Further, from DE 199 57 371 an aqueous solution of mitomycin C is known, which comprises buffer and has a pH in the range from 6.0 to 9.0. In addition, the solution is characterized by a specific ratio of mitomycin C concentration to buffer ionic strength.

30 However, the shelf life of ready-to-use solutions of mitomycin is not sufficient, since the active substance content decreases markedly on storage. For this reason, mitomycin C is

prepared in the form of lyophilisates or dry powders, and these pharmaceutical forms are reconstituted with sterile solvent directly before administration, so as to give ready-to-use solutions. Lyophilisates are marketed for example under the brand name "Mitem". Dry powders, for example mixed with NaCl, are for example obtainable under the brand name "Mito-medac".

5

10

However, the lyophilisates available on the market have the disadvantage that only a relatively short shelf life of at most two years is stated for them. This is due to the high instability of mitomycin C, which readily decomposes in the lyophilisation solution with formation of a range of degradation products. The main degradation products are albomitomycin C, D1 and D2, the structural formulae of which are reproduced below together with that of mitomycin C.

Mitomycin C

#### Albomitomycin C



D1



Although dry powders of mitomycin C have a longer shelf life of about four years, they have the disadvantage that during their preparation dusts are formed which result in health-endangering

D2

contamination of the personnel and contamination of the production plants. The long period of time necessary for complete dissolution of these dry powders is also not satisfactory.

With the known lyophilisates and dry powders, the reconstitution to solutions is usually effected to give end concentrations of 1 mg/ml mitomycin C, e.g. 20 mg mitomycin C to 20 ml water or isotonic saline solution (0.9% wt.-%). Lower concentrations after dilution to 0.2 and 0.15 mg/ml for ophthalmological applications are also known. However, because of the poor solubility of the known products, a higher end concentration in water is not possible. In the literature, their saturation concentration is stated as 0.5-1.0 mg/ml.

10

15

The object of the invention is therefore to avoid the disadvantages of the known lyophilisates and dry powders, and in particular to provide a process for the preparation of a freeze-dried pharmaceutical composition of mitomycin C, which process does not result in a significant reduction in the active substance content and which results in a composition which has high stability and purity and hence exhibits only slight degradation of the active substance even on prolonged storage. In addition, the pharmaceutical composition produced should be rapidly and completely reconstitutable to form ready-to-use solutions which can have a high concentration of active substance.

20 This object is achieved by the freeze-dried pharmaceutical composition according to claims 1 to 8.

The freeze-dried pharmaceutical composition according to the invention comprising mitomycin C is characterized in that it comprises urea and is obtainable by a process, in which a solution of mitomycin C is freeze-dried, wherein the solution comprises at least one organic solvent.

The solution subjected to the freeze-drying in the process is also referred to below as "lyophilisation solution".

The organic solvent used in the process is in particular selected from the group comprising tert.butanol, n-propanol, n-butanol, isopropanol, ethanol, methanol, 1-pentanol, chlorobutanol, acetic acid, acetone, dimethyl carbonate, acetonitrile, dichloromethane, methyl ethyl ketone, methyl isobutyl ketone, methyl acetate, carbon tetrachloride, dimethyl sulphoxide, N,Ndimethylacetamide, hexafluoroacetone, dimethyl sulphone and cyclohexane. The organic solvent is preferably selected from the group of tert.-butanol, ethanol, isopropanol, acetone, dimethyl sulphoxide and N,N-dimethylacetamide. Particularly preferably, the organic solvent is

5

20

tert.-butanol.

10 In one embodiment, the organic solvent is the only solvent in the solution. When organic solvents which are solid at room temperature are used, the mitomycin C is usually dissolved in previously melted organic solvent.

In a further, preferred embodiment, the solution used in the process according to the invention 15 also comprises water.

The dissolution of mitomycin C in at least one organic solvent, preferably in a mixture with water, surprisingly results in a lyophilisation solution which is physically and chemically stable at room temperature for up to 48 h and at 5°C even up to 9 days. At the same time, the content of impurities in this solution also remains surprisingly low and thus the purity of the solution remains very high. Because of the high stability and the high purity of the solution, very pure freeze-dried compositions of mitomycin C can be prepared with the process.

The freeze-drying of this solution surprisingly results in no significant increase in impurities, so that the freeze-dried composition obtained also exhibits high purity. The composition also surprisingly exhibits very high stability and even after storage for several months it exhibits a high content of active substance and only small quantities of undesired degradation products.

Further, the composition produced with the process, also referred to below as "lyophilisate",
exhibits a surprisingly high solubility in the solvents usually used for reconstitution and enables the preparation of ready-to-employed solutions which comprise mitomycin C at a concentration

of more than 1 mg/ml. Hence, by using the composition, ready-to-use solutions with a particularly high active substance concentration can be produced.

5

The solution used in the process preferably comprises a mixture of ethanol and water, a mixture of isopropanol and water, a mixture of acetone and water, a mixture of dimethyl sulphoxide and water, or a mixture of dimethyl sulphoxide and dimethylacetamide. Particularly preferably the solution comprises a mixture of tert.-butanol and water.

The mixture of ethanol and water used in particular comprises at least 1 wt.-%, preferably at least
5 wt.-%, particularly preferably at least 10 wt.-% and quite particularly preferably at least 15 wt.-%
% ethanol.

The mixture of isopropanol and water used in particular comprises at least 1 wt.-%, preferably at least 5 wt.-%, particularly preferably at least 10 wt.-% and quite particularly preferably at least 15 wt.-% isopropanol.

The mixture of acetone and water used in particular comprises at least 1 wt.-%, preferably at least 5 wt.-%, particularly preferably at least 10 wt.-% and quite particularly preferably at least 15 wt.-% acetone.

20

15

5

The mixture of dimethyl sulphoxide and water used in particular comprises at least 1 wt.-%, preferably at least 30 wt.-%, particularly preferably at least 50 wt.-% and quite particularly preferably at least 80 wt.-% dimethyl sulphoxide.

25 The mixture of dimethyl sulphoxide and dimethylacetamide used in particular comprises at least 5 wt.-% and preferably at least 50 wt.-% dimethyl sulphoxide.

In the process, a solution which comprises a mixture of tert.-butanol and water is particularly preferably used. This mixture of tert.-butanol and water in particular comprises at least 1 wt.-%,

preferably at least 5 wt.-%, particularly preferably at least 10 wt.-% and more preferably at least

30

15 wt.-% tert.-butanol.

In a quite particularly preferred embodiment, the mixture of tert.-butanol and water used comprises 80 to 99 wt.-%, in particular 84 to 95 wt.-%, preferably 88 to 92 wt.-% and quite particularly preferably about 89 wt.-% tert.-butanol. In an alternative quite particularly preferred embodiment, the mixture of tert.-butanol and water used comprises 10 to 30 wt.-%, in particular 15 to 25 wt.-% preferably 18 to 22% and quite particularly preferably about 20 wt.-% tert.-

5 15 to 25 wt.-%, preferably 18 to 22% and quite particularly preferably about 20 wt.-% tert.butanol.

The lyophilisation solution used in the process usually comprises mitomycin C at a concentration of 0.1 to 500 mg/g, in particular 0.5 to 10.0 mg/g, preferably 0.8 to 6.0 mg/g and particularly preferably 1.0 to 4.0 mg/g.

The solution also comprises urea. The use of this additive inert towards the active substance results in a lyophilisate which, because of its special properties, protects the active substance against destabilizing influences and thus preserves the active substance in its pure form.

15

10

In a preferred embodiment of the process, the solution comprises urea at a concentration of 1 to 200 mg/g, in particular 5 to 100 mg/g and preferably 10 to 60 mg/g.

The solution can also comprise polyethylene glycol or mannitol.

20

The polyethylene glycol has in particular an average molecular weight (number average) of 1000 to 8000 and preferably of 2000 to 8000. Particularly preferably, PEG 4000, which has an average molecular weight (number average) of about 4000, is used as polyethylene glycol.

In a preferred embodiment of the process, the solution comprises polyethylene glycol at a concentration of 1 to 200 mg/g, in particular 5 to 100 mg/g and preferably 10 to 60 mg/g.

In a preferred embodiment of the process, the solution comprises mannitol at a concentration of 1 to 200 mg/g, in particular 5 to 100 mg/g and preferably 10 to 60 mg/g.

The solution can additionally also comprise other usual additives such as further sugar alcohols, e.g. isomalt, lactitol, sorbitol, lactitol, sorbitol, xylitol, threitol, erythritol or arabitol, sugars, e.g. sucrose, glucose, fructose, maltose, rhamnose, lactose or trehalose, amino acids, e.g. L-phenylalanine, L-tryptophan, L-proline, L-histidine, L-glycine or L-arginine, polymers, e.g. polyvinylpyrrolidones, polyvinyl acetates, starch derivatives, e.g. cyclodextrins, dextroses, amylopectins, amyloses, polysaccharides, e.g. alginates, pectins, celluloses, agents for adjusting isotonicity, e.g. sodium chloride, calcium chloride dihydrate, sodium acetate trihydrate, disodium hydrogen phosphate dihydrate, and/or agents for adjusting the pH, e.g. trometamol, usually at concentrations in each case of 1 to 200, in particular 5 to 100 and preferably 10 to 60 mg/g.

The freeze-drying of the lyophilisation solution is as a rule effected using freeze-drying processes and machines normally used for pharmaceutical purposes. As a rule, the solution is filled into a suitable vessel, and the vessel is placed in a conventional freeze-dryer with coolable and heatable surfaces on which the solution can be exposed to the various temperatures of the freeze-drying process. To achieve the drying, the solution is usually frozen and exposed to a decreased atmospheric pressure. As a result, sublimation of the solvent from the frozen solution takes place to a great extent, which precipitates for example on cooler regions of the freeze-dryer provided for this. This is then as a rule also followed by a secondary drying at higher temperatures. After completion of the freeze-drying, the lyophilisate obtained is as a rule allowed to come to room temperature and the vessel used is sealed under sterile conditions. A suitable programme for the lyophilisation can comprise charging of the vessel at room temperature, freezing at -50°C to -35°C under normal pressure, subsequent lowering of the atmospheric pressure and then increasing of the temperature by 25 to 95°C in order to effect the drying.

25

30

5

10

The freeze-dried composition obtained with the process evidently also has a special and advantageous structure, since only a very short period of less than 60 seconds, in particular less than 30 seconds and preferably 4 to 20 seconds suffices for complete reconstitution to form a ready-to-use solution. This is an important advantage, since as a result it is possible for clinic staff to prepare ready-to-use solutions freshly directly before the intended administration to patients, without having to allow for long waiting times for complete dissolution. Likewise, with

such short reconstitution times, the risk of undesired degradation reactions of the active substance decreases.

For the reconstitution of the composition, isotonic saline solution or water for injection is usually employed. Other pharmaceutically acceptable solutions are also possible for the reconstitution, such as for example Ringer's lactate solutions or phosphate buffers.

The freeze-dried pharmaceutical composition according to the invention comprises urea. Such a composition requires smaller quantities of reconstituting agent for reconstitution and thus a

10 ready-to-use solution with a particularly high concentration of mitomycin C can be prepared by means thereof.

The composition usually comprises urea at a concentration of 0.2 to 1, in particular 0.5 to 0.99, preferably 0.8 to 0.95 and particularly preferably 0.93 to 0.94 g per gram of composition.

15

5

The composition can also comprise polyethylene glycol usually at a concentration of 0.2 to 1, in particular 0.5 to 0.99, preferably 0.8 to 0.95 and particularly preferably 0.93 to 0.94 g per gram of composition.

- 20 The composition can also comprise mannitol usually at a concentration of 0.2 to 1, in particular 0.5 to 0.99, preferably 0.8 to 0.95 and particularly preferably 0.93 to 0.94 g per gram of composition.
- The composition according to the invention comprises the impurities D1, D2 and albomitomycin C in a total quantity of in particular less than 2.0%, preferably less than 1.5%, particularly preferably less than 1.0%, more preferably less than 0.8% and quite particularly preferably less than 0.6%, wherein the quantity of these impurities is determined according to the impurities process according to Ph Eur 8.0 Mitomycin Monograph 01/2008: 1655. In the examples, this determination process is also referred to as process 4.

The invention is explained in more detail below on the basis of examples.

**Examples** (Examples in accordance with the claims comprise urea and are obtainable by a process, in which a solution of mitomycin C is freeze-dried, wherein the solution comprises at least one organic solvent.)

#### **Example 1 – Solubility and stability of mitomycin C in various solvents**

The solubility and stability of mitomycin C in the various solvents and solvent mixtures stated 10 in Table 1 were investigated.

No.		Cosolvent wt	Water wt	Mitomycin C	Mannitol
		%	%	concentration[mg/g	concentration [mg/g]
1	Water	-	100	2.0	4.0
2	Tertbutanol	32.6	67.4	2.1	4.1
3	Ethanol	16.4	83.6	2.2	4.4
4	Isopropanol	16.2	83.8	2.2	4.3
5	Acetone	7.1	92.9	2.0	4.1

### Table 1 "Solvents/solvent mixtures used"

15

5

For this purpose, solutions which contained 2.0 to 2.2 mg/g mitomycin C and 4.0 to 4.4 mg/g mannitol were prepared using each of these mixtures.

20

After 1.5 hours' stirring time at room temperature and with exclusion of light, the mixtures obtained were examined for particles under irradiation with linearly polarized light. The result of the visual examination is presented in Table 2.

No.	Solvent/solvent mixture	Description of the mixtures	
	wt%		
1	Water 100%	dark blue suspension, not completely	
		dissolved	
2	Tertbutanol 32.6%	dark blue clear solution, completely	
		dissolved	
3	Ethanol 16.4%	dark blue clear solution, completely	
		dissolved	
4	Isopropanol 16.2%	dark blue clear solution, completely	
		dissolved	
5	Acetone 7.1%	dark blue clear solution, completely	
		dissolved	

### Table 2 "Visual examination"

Further, the content of mitomycin C and impurities in the mixtures obtained was determined
after 24 hours' storage at room temperature. The corresponding results are presented in Table
3.

No.	Solvent mixture wt%	Mitomycin C	Total content of impurities
		content [%]*	(D1, D2 and albomitomycin
			C) [%]**
2	Tertbutanol 32.6%	98.04	1.96
3	Ethanol 16.4%	94.74	5.26
4	Isopropanol 16.2%	95.69	4.30
5	Acetone 7.1%	93.99	6.01

### Table 3 "Content of mitomycin C and impurities after 24 h storage at room temperature"

5 \* Determination by process 1 (see Table 5)

\*\* Determination by process 2 (see Table 5)

The results presented in Table 3 show that the mitomycin C in the lyophilisation solution in a mixture of tert.-butanol and water used according to the invention has particularly high stability.

10

Further, the mixtures obtained were again examined for visible particles under irradiation with linearly polarized light after 24 hours' storage with exclusion of light at (a) room temperature and (b) 2 to 8°C. The results thus obtained are presented in Table 4.

No.	Solvent/solvent mixture	Storage at room	Storage at 2 to 8°C
	wt%	temperature	
1	Water 100%	dark blue suspension;	dark blue suspension
		dispersed particles	
2	Tertbutanol 32.6%	dark blue clear solution	dark blue clear solution
3	Ethanol 16.4%	dark blue clear solution;	dark blue clear solution;
		needle-like particles	needle-like particles
4	Isopropanol 16.1%	dark blue clear solution	dark blue clear solution;
			needle-like particles
5	Acetone 7.1%	dark blue clear solution;	dark blue clear solution;
		needle-like particles	needle-like particles

#### Table 4 "Visual examination after storage for 24 h"

5 With the solvent mixtures usable according to the invention, markedly more stable solutions were obtained than with water. The solution obtained with the mixture of 32.6 wt.-% tert.-butanol and 67.4 wt.-% water proved to be the most stable, since no particles were visible in it even after 24 h both at room temperature and also at 2 to 8°C. Furthermore, in the tert-butanol-water mixture after 24 hours' storage at room temperature the content of mitomycin C was the highest and the content of impurities the lowest.

### Determination of the content of mitomycin C and of impurities by high pressure liquid chromatography

15

The samples are each diluted to 0.5 mg/ml mitomycin C with N,N-dimethylacetamide. The reference solution contains 0.5 mg/ml mitomycin C in N,N-dimethylacetamide.

	Process 1	Process 2
Detection:	UV at 365 nm	UV at 254 nm
Mobile	Buffer/methanol 62.5/37.5, vol/vol	Buffer/methanol
phase:		75/25, vol/vol
	Buffer:	
	0.025% acetic acid in 0.02 M	Buffer:
	ammonium acetate	0.025% acetic acid in 0.02 M
		ammonium acetate
Flow rate:	1.4 ml/min	2.6 ml/min

### Table 5 "High pressure liquid chromatography conditions"

Column:

YMC-Pack Phenyl 4.6 mm x 300 mm, 10 µm

5 Temperature: 25°C Injection volume: 10 μl

# Example 2 – Solubility and stability of mitomycin C in dimethyl sulphoxide and a 10 mixture of dimethyl sulphoxide and N,N-dimethylacetamide

The solubility and stability of mitomycin C in the solvents stated in Table 6, namely in dimethyl sulphoxide (DMSO) and a mixture of dimethyl sulphoxide (DMSO) and N,N-dimethyl-acetamide (DMAA) were investigated.

15

### Table 6 "DMSO or DMSO/DMAA as solvent/solvent mixture"

No.	Dimethyl sulphoxide wt%	N,N-Dimethylacetamide wt%
6	5.7	94.3
7	100	-

For this purpose, solutions using each of these solvents were prepared, which for No. 6 (DMSO/DMAA) contained 0.53 mg/g mitomycin C and 8.0 mg/g urea and for No. 7 (DMSO) 1.82 mg/g mitomycin C and 27.3 mg/g urea. The solutions were stored at room temperature and with exclusion of light for a period of up to 71.5 hours.

Table 7a "Content of mitomycin in DMSO/DMAA mixture"

5

		Mitomycin content [%]				
DMSO/DMAA	t=0	t=6.5h	t=14.5h	t=21h	t=42h	t=71.5h
wt%						
5.7/94.3 1	100	99.93	99.62	98.84	99.11	98.58

<sup>1</sup> Content determined with process 3

	Mitomycin purity [%]					
DMSO wt%	t=0	t=4h	t=6.5h	t=13h	t=20h	t=26.5h
100 <sup>2</sup>	100	99.31	99.32	99.19	99.36	99.32

### Table 7b "Purity of mitomycin in DMSO"

<sup>5</sup> <sup>2</sup> Purity determined with process 4; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

In both solvents, the mitomycin C proved to be very stable. The use of DMSO alone resulted in a particularly small decrease in the original content of mitomycin C from 100% to 99.32% within a period of 26.5 hours.

10

15

The content of mitomycin C and hence the stability of the solutions obtained was checked by processs 3 and 4. Process 3 corresponds to the content process according to Ph Eur 8.0 Mitomycin Monograph 01/2008: 1655. Process 4 corresponds to the impurities process according to Ph Eur 8.0 Mitomycin Monograph 01/2008: 1655.

### Example 3 – Solubility and stability of mitomycin C in various tert.-butanol/water mixtures

5 The solubility and stability of mitomycin C in the various mixtures of tert.-butanol and water stated in Table 8 were investigated.

For this purpose, solutions using each of these mixtures were prepared, wherein the concentration of mitomycin C was 2.1 mg/g for mixture No. 8, 2.2 mg/g for mixture No. 9 and 2.4 mg/g for mixture No. 10.

These solutions were stored for 16 h with exclusion of light at (a) room temperature and (b) at 2 to 8°C and then again examined for visible particles under irradiation with linearly polarized light. The results thus obtained are also presented in Table 8.

15

10

No.	Solvent mixture wt% Storage at room		Storage at 2 to 8°C (b)	
		temperature (a)		
8	tertbutanol 32.6%	dark blue clear solution	dark blue clear solution	
9	tertbutanol 51.7%	dark blue clear solution	dark blue clear solution	
10	tertbutanol 72.9%	dark blue clear solution	dark blue clear solution	

Table 8 "Visua	l examination	after storage	for 16 h"
----------------	---------------	---------------	-----------

20

All mixtures of tert.-butanol and water were capable of dissolving the stated quantity of mitomycin C, and in the solutions obtained no particles were visible even after 16 h storage.

25

For the HPLC determination of the stability of mitomycin C in tert.-butanol/water mixtures, mitomycin C was added to a mixture of 20 wt.-% tert.-butanol and 80 wt.-% water (No. 11) in a quantity such as to obtain a solution with a concentration of 10 mg/g mitomycin C. Further,

mitomycin C was added to a mixture of 32.6 wt.-% tert.-butanol and 67.4 wt.-% water (No. 12) in a quantity such as to obtain a solution with a concentration of 2.1 mg/g mitomycin C. Further, mitomycin C was added to a mixture of 89 wt.-% tert.-butanol and 11 wt.-% water (No. 13) in a quantity such as to obtain a solution with a concentration of 5 mg/g mitomycin

- 5 C. Further, a mitomycin-compriseing solution in pure tert.-butanol (100 wt.-%) (No. 14) and for comparison a solution in water (No. 15) were prepared. In the experiment with pure tert.-butanol, this was firstly melted by warming to about 30°C and the mitomycin C was then dissolved in the melt.
- 10 The solutions were then stored for 24 h at room temperature in brown glass vessels. The respective results are presented in Table 9.

No.	Solvent mixture wt	Mitomycin	Mitomycin	Mitomycin	Mitomycin
	%	[mg/g]	purity*	purity* t =	purity* t=
			t = 0h [%]	6h [%]	24h [%]
11	Tertbutanol 20%	10	99.51	97.01	93.55
12	Tertbutanol 32.6%	2.1	100.0	99.4	98.7
13	Tertbutanol 89%	5	99.56	98.95	98.47
14	Tertbutanol 100%	0.38	100.0	100.0	100.0
15	Water (Reference)	0.48	96.8	92.1	89.9

Table 9	"Stability	of mitomyci	n C in tert.	butanol/wate	r mixtures"
---------	------------	-------------	--------------	--------------	-------------

5

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

In the tert.-butanol-containing solutions, the mitomycin C had high stability, wherein the use of the mixtures No. 12 and No. 13 resulted in a particularly slight decrease in the original purity at t=0 of mitomycin C from 100.0% to 98.7% and 99.56% to 98.47%. A particularly high purity of 100.0 wt.-% after 24 h could be achieved in 100 wt.-% tert.-butanol, however, the solubility of mitomycin C in pure tert.-butanol is lower.

15 In contrast to this, the purity of mitomycin C in water after 24 h is significantly worse (at t=0 from 96.8% to 89.9%).

### Example 4 - Compatibility of further additives with tert.-butanol/water mixtures

20

Several solutions were prepared, each by dissolving 20 mg mitomycin C in 6 g of a mixture of 89 wt.-% tert. butanol and 11 wt.-% water. The additives urea, PEG 4000 or trometamol were

added to these solutions, in order in each case to give a concentration of 50 mg/g. In addition, the solution with trometamol was adjusted to a pH of 7.4 with acetic acid.

19

At t=0 and after 6 h and 24 h at room temperature, the solutions were examined for their 5 content of mitomycin C and compared with a solution which contained no further additive. The determination of the content was performed according to process 2, and the results are presented in Table 10.

# Table 10 "Stability of solutions with urea,polyethylene glycol and trometamol"

Additive	Mitomycin C	Mitomycin C	Mitomycin C
	purity*	purity*	purity*
	t=0	t=6 h	t=24 h
Urea	99.69	99.76	99.14
PEG 4000	100	99.58	99.50
Trometamol	99.56	99.44	99.50
Without additive	99.61	99.63	99.48

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

15

### Example 5 - Stability of mitomycin C in urea-containing tert. butanol/water mixtures

20

Urea was dissolved in a mixture of 89 wt.-% tert.-butanol and 11 wt.-% water in a quantity such as to achieve a concentration of 5 wt.-%. Next, mitomycin C was added in a quantity such as to obtain a concentration of 3.33 mg/g, and the solution was stirred for one hour. The solution was filled into brown glass vessels and stored for 9 days at room temperature (RT) without protection against light, at room temperature (RT) with exclusion of light and at 5°C with exclusion of light.

Purity determinations according to process 2 were performed at different times. The results are presented in Table 11.

5

### Table 11 "Stability of mitomycin solutions with urea"

		Mitomycin purity* [%]				
Storage	t=0	t=6h	t=24h	t=48h	t=72h	t=9d
RT; without protection	99.32	98.83	97.60	96.77	96.14	95.78
against light						
RT; with exclusion of	99.33	98.83	97.80	97.04	96.74	96.35
light						
5°C; with exclusion of	99.23	99.18	99.20	98.86	98.80	98.21
light						

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

10

20

Even after nine days' storage, under all conditions, the solution was still surprisingly stable and the purity of the solution surprisingly high.

### 15 <u>Example 6 – Freeze-drying of solutions of mitomycin C in various tert.-</u> <u>butanol/water mixtures</u>

Firstly, solutions of mitomycin C in mixtures of tert.-butanol and water with 16.1 wt.-%, 32.6 wt.-% and 72.9 wt.-% tert.-butanol were prepared. These solutions contained 2.1 mg/g mitomycin C and 4.1 mg/g mannitol for the solution with the mixture with 16.1 wt.-% tert.-butanol, 2.2 mg/g mitomycin C and 4.3 mg/g mannitol for the solution with the mixture with 32.6 wt.-% tert.-butanol and 2.4 mg/g mitomycin C and 4.8 mg/g mannitol for the solution with the mixture with 72.9 wt.-% tert.-butanol.

All solutions were clear and sufficiently stable.

For comparison, a solution of mitomycin C in water, which contained 7 mg/g mannitol and because of the poor solubility in water 0.7 mg/g mitomycin C, was also prepared.

By conventional freeze-drying with (i) freezing of the solutions, (ii) application of a vacuum, (iii) primary and (iv) secondary drying, lyophilisates were obtained. These were reconstituted with water, and the reconstituted solutions obtained were analysed according to Example 1 for their content of mitomycin C and of impurities. The results are presented in Table 12.

## Table 12"Content of mitomycin C and impurities after freeze-drying and<br/>reconstitution"

Quantity of tertbutanol in	Mitomycin C	Impurities Content*** [%]
mixture wt%	Content** [%]	Total; D1; D2; albomitomycin C
Tertbutanol 0%	98.36	1.64; 0.40; 0.48; 0.76
Tertbutanol 16.1%	99.55	0.46; ND*; 0.15; 0.31
Tertbutanol 32.6%	99.81	0.20; ND*; 0.10; 0.10
Tertbutanol 72.9%	100.00	ND*; ND*; ND*; ND*

15

25

5

10

\* ND = not detectable, i.e. below the detection limit\*\* HPLC determination by process 1

\*\*\* HPLC determination by process 2

20 After lyophilisation and reconstitution, all mixtures of tert.-butanol and water used according to the invention resulted in solutions with a very high content of mitomycin C and only a very low content of impurities (D1, D2 and albomitomycin C).

In contrast to this, lyophilisation using water gave a reconstituted solution with a markedly lower content of mitomycin C and a very significant quantity of impurities.

### Example 7 Stability of mitomycin C in tert.-butanol-water mixture (89/11) with 20 mg/g urea

5

10

20

Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol and 11 wt.-% water were prepared. As well as 1.33 mg/g mitomycin C, the solutions also each contained 20 mg/g urea. At t=0 and after 5, 22 and 27 hours at room temperature (RT) with exclusion of light and in the refrigerator, the purity of the solutions was investigated according to process 2. The results are shown in Table 13.

### Table 13 "Stability of mitomycin C in 20 mg/g urea-containing solutions"

	Mitomycin C purity* [%]		
Conditions	t=5h	t=22h	t=27h
RT; with exclusion of light	99.7	99.2	98.9
5°C; with exclusion of light	99.8	99.8	99.7

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

Even after more than one day's storage at room temperature, the purity of the solution is surprisingly high, which reflects the surprisingly high stability of the active substance in the solvent.

### Example 8 - Stability of mitomycin C in tert.-butanol- water mixture (89/11) with 25 mg/g urea

25 Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol and 11 wt.-% water were prepared. As well as 1.67 mg/g mitomycin C, the solutions also each contained 25 mg/g urea.

At t=0 and after 5, 22 and 27 hours at room temperature (RT) with exclusion of light and in the refrigerator, the purity of the solutions was investigated according to process 2.

Table 14 "Stability of mitomycin C in 25 mg/g urea-containing solutions"

5

	Mitomycin C purity* [%]		
Conditions	t=5h	t=22h	t=27h
RT; with exclusion of light	99.6	99.1	98.6
5°C; with exclusion of light	99.8	99.5	99.5

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

10 Even after more than one day's storage at room temperature, the purity of the solution is surprisingly high, which reflects the surprisingly high stability of the active substance in the solvent.

# Example 9 – Freeze-drying of solutions of mitomycin C containing urea or polyethylene glycol (50 mg/g)

5 Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol and 11 wt.-% water were prepared. As well as 3.33 mg/g mitomycin C, the solutions also each contained either 50 mg/g urea or PEG 4000. In each case, 6 g of these solutions was filled into small vials and freezedried in conventional manner. The lyophilisate cakes containing urea and PEG 4000 were firm and defect-free.

10

The lyophilisates were analysed for the purity of mitomycin C directly after their preparation (t = 0) and after storage for 1 month (t = 1 month) at 40°C. The results are presented in Table 15.

15

### Table 15 "Stability of lyophilisates with urea and polyethylene glycol"

t = 0;		
Additive	Mitomycin purity *	Reconstitution time
	[%]	
Urea	99.65	<< 30 secs
PEG 4000	99.34	<< 30 secs
t = 1 month at 40	<b>)°С;</b>	
Additive	Mitomycin purity *	Reconstitution time
	[%]	
Urea	99.89	<< 30 secs
PEG 4000	96.19	<< 30 secs

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

Even after storage at an elevated temperature of 40°C, the lyophilisates produced had a high purity of mitomycin C, which reflects their very good stability.

Moreover, the lyophilisates could be completely reconstituted with isotonic saline solution in very much less than 30 secs.

5

#### **Example 10 – Freeze-drying of solutions of mitomycin C containing urea (20 mg/g)**

- Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol (TBA) and 11 wt.-% water were prepared. As well as 1.33 mg/g mitomycin C, the solutions also each contained 20 mg/g urea. In each case, 1.5 g of this solution was filled into small vials and freeze-dried in conventional manner. The lyophilisate cakes were firm and defect-free.
- 15 The lyophilisates were analysed for the purity of mitomycin C directly after their preparation (t=0). In addition, the reconstitution time was determined. The reconstitution was effected with isotonic saline solution to an end concentration of 1 mg/g mitomycin C. The results are presented in Table 16.

### Table 16

### "Characteristics of urea-containing lyophilisates"

$\mathbf{t} = 0$		
Additive Mitomycin C purity*		Reconstitution time
	[%]	
Urea	99.5	<< 30 secs

5

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

The lyophilisates produced had a high purity of mitomycin C, which reflects the very good stability of the active substance during the process.

Moreover, the lyophilisates could be completely reconstituted with isotonic saline solution in very much less than 30 secs.

### 15 <u>Example 11 – Freeze-drying of solutions of mitomycin C containing urea (25 mg/g)</u>

Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol (TBA) and 11 wt.-% water were prepared. As well as 1.67 mg/g mitomycin C, the solutions also each contained 25 mg/g urea. In each case, 1.2 g of this solution was filled into small vials and freeze-dried in

20 conventional manner. The lyophilisate cakes with urea were firm and defect-free.

## Table 17"Urea-containing lyophilisates"

t = 0		
Additive	Mitomycin purity	Reconstitution time
	[%]*	
Urea	99.5	<< 30 secs

5

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

10

The lyophilisates produced had a high purity of mitomycin C, which reflects the very good stability of the active substance during the process. Moreover, the lyophilisates could be completely reconstituted with isotonic saline solution in very much less than 30 secs.

### Example 12 - Mitomycin C in tert.-butanol/water mixture (95/5) with mannitol (45 mg/g)

Solutions of mitomycin C in a mixture of 95 wt.-% tert.-butanol (TBA) and 5 wt.-% water were prepared. As well as 5 mg/g mitomycin C, the solutions also each contained 45 mg/g mannitol. In each case, 4.0 g of this solution was filled into small vials and freeze-dried in conventional manner. The lyophilisate cakes were firm and defect-free.

### Table 18"Mannitol-containing lyophilisate"

28

t = 0		
Additive	Mitomycin purity*	
	[%]*	Reconstitution time
Mannitol	99.6	<< 30 secs

5

\* HPLC determination by process 2; purity is defined as area of the mitomycin peak relative to the total area of all peaks in the HPLC chromatogram

10

The lyophilisates produced had a high purity of mitomycin C, which reflects the very good stability of the active substance during the process. Moreover, the lyophilisates could be completely reconstituted with isotonic saline solution in very much less than 30 secs.

### Example 13 - Stability of mitomycin lyophilisates

15

20

25

Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol (TBA) and 11 wt.-% water were prepared. As well as 3.33 mg/g mitomycin C, the solutions also each contained 50 mg/g urea. In each case, 6 [corresponds to 20 mg mitomycin C] or 12 g [corresponds to 40 mg mitomycin C] of these solutions was filled into small vials and freeze-dried in conventional manner. The lyophilisates obtained were examined at different times t=0, t=3 months and t=6 months at 40°C and 75% relative humidity (r.h.).

The determination of the content of mitomycin C was performed by process 3 and that of the impurities content by process 4. The reconstitution was effected with 20 and 40 ml respectively of isotonic saline solution to an end concentration of 1 mg/ml.

### Table 19

### "Stability at 40°C and 75% r.h. [20 mg mitomycin C]"

	t=0	t=3 months	t=6 months
Reconstitution time	7 secs	12 secs	17 secs
Mitomycin C content*	97.8%	98.2%	98.3%
Impurity	0.29%	0.20%	0.26%
albomitomycin C**			
Sum of all impurities**	0.29%	0.55%	0.61%

5 \* determination by process 3 \*\* determination by process 4

#### Table 20

### "Stability at 40°C and 75% r.h. [40 mg mitomycin C]"

1	0

	t=0	t=3 months	t=6 months
Reconstitution time	6 secs	20 secs	18 secs
Mitomycin C content*	97.2%	96.6%	96.7%
Impurity	0.32%	0.22%	0.21%
albomitomycin C**			
Sum of all impurities**	0.32	0.55	0.58

\* determination by process 3

\*\* determination by process 4

15 Even after 6 months' storage at elevated temperature of 40°C and elevated atmospheric humidity of 75% r.h., the lyophilisates produced contained only very small quantities of impurities. Moreover, the lyophilisates could be completely reconstituted with isotonic saline solution in very much less than 30 secs.

### 20 Example 14 - Elevated end concentration after reconstitution

Solutions of mitomycin C in a mixture of 89 wt.-% tert.-butanol and 11 wt.-% water were prepared. As well as 3.33 mg/g mitomycin C, the solutions also each contained 50 mg/g urea. In each case, 6 g [corresponds to 20 mg mitomycin C] of these solutions was filled into small vials and freeze-dried in conventional manner. The lyophilisates obtained were reconstituted

5 with 20 ml or 10 ml water for injection.

Mitomycin C	Mitomycin C content*	Description	of	the
end concentration	[%]	reconstituted solution		
1 mg/ml	100.6	Clear, free	e from	visible
		particles		
2 mg/ml	100.6	Clear, free	e from	visible
		particles		

\* determination by process 3

The reconstituted solutions in water for injection had a high content of mitomycin C. The 10 solutions were clear and free from visible particles.

### Patentkrav

1. Frysetørket farmasøytisk sammensetning med et innhold av mitomycin C som inneholder urea og kan oppnås ved en fremgangsmåte der en løsning av mitomycin C frysetørkes, hvor løsningen inneholder minst ett organisk løsemiddel.

**2.** Sammensetningen ifølge krav 1, som inneholder minst ett hjelpestoff valgt fra gruppen av polyetylenglykol og mannitol.

- 3. Sammensetningen ifølge krav 1 eller 2, som inneholder urea i en konsentrasjon på 0,2 til 1, spesielt 0,5 til 0,99, foretrukket 0,8 til 0,95 og spesielt foretrukket 0,93 til 0,94 g per gram av sammensetningen.
- 4. Sammensetningen ifølge krav 2 eller 3, som inneholder polyetylenglykol i en
   konsentrasjon på 0,2 til 1, spesielt 0,5 til 0,99, foretrukket 0,8 til 0,95 og spesielt
   foretrukket 0,93 til 0,94 g per gram av sammensetningen.

5. Sammensetningen ifølge et av kravene 2 til 4, som inneholder mannitol i en konsentrasjon på 0,2 til 1, spesielt 0,5 til 0,99, foretrukket 0,8 til 0,95 og spesielt foretrukket 0,93 til 0,94 g per gram av sammensetningen.

**6.** Sammensetningen ifølge et av kravene 2 til 5, der polyetylenglykol har et gjennomsnittlig molekylvekt (antallsgjennomsnitt) på 1000 til 8000 og foretrukket 2000 til 8000.

25

20

5

**7.** Sammensetning ifølge et av kravene 1 til 6, som inneholder forurensningene D1, D2 og albomitomycin C i en total lengde på mindre enn 2,0 %, foretrukket mindre enn 1,5 %, spesielt foretrukket mindre enn 1,0 %, ytterligere foretrukket mindre enn 0,8 % og helt spesielt foretrukket mindre enn 0,6 %, hvor mengden av disse forurensningene bestemmes ifølge forurensningsmetoden ifølge Ph Eur 8.0 mitomycin monografi 01/2008: 1655.

**8.** Sammensetning ifølge ett av kravene 1 til 7, hvor løsningen dessuten inneholder vann.

35